




# UCLPartners Proactive Care Framework: Managing Hypertension & Cardiovascular risk

Version 9

**The Proactive Care Frameworks (PCF) can be used independently or in conjunction with [CVD ACTION](#)**

**To request the PCF search tools please visit:  
[UCLPartners Proactive Care Frameworks - UCLPartners](#)**

# Background to the Frameworks



# The Challenge of Long-Term Condition Management in Primary Care

## Historical challenge in long term condition care

- Late diagnosis, suboptimal treatment, unwarranted variation
- Focus on reactive rather than proactive care
- Limited support for self-management



## Real World Primary Care

- Complexity, multimorbidity and time pressures
- Often asymptomatic conditions
- Soaring demand and shifting priorities



## This is a wicked problem

- Despite decades of QOF incentives, NICE guidance and myriad quality schemes, improvement has been marginal at best
- Primary care needs support to do things differently and at scale



# UCLPartners Proactive Care Frameworks Address Core Challenges in Primary Care

## Aim

Help people with long term conditions to stay well longer

## Objectives

1. Mobilise data - Identify patients whose care needs optimising and prioritise those at highest risk
2. Harness wider workforce - standardise delivery of holistic proactive care by wider primary care team
3. Support GPs to safely manage workflow, improve care and outcomes by releasing capacity

## Framework components

- ✓ Risk stratification & prioritisation tools
- ✓ Locally adaptable resources to support real world management
- ✓ Systematic use of wider primary care team (eg ARRS\* roles) to deliver structured support for education, self-management and behaviour change

## Framework Development

- Led by primary care clinicians
- Based on NICE guidelines and clinical consensus
- Patient and public support



# Why Hypertension? The Case for Change

# Why Focus on Hypertension and Cardiovascular Risk?

- Hypertension is the leading risk factor causing death worldwide
- In England, there are:
  - An estimated 1.8 million people with undetected hypertension
  - 2.9 million adults with diagnosed hypertension who are not achieving the age-appropriate BP treatment target
- Delaying intervention for more than 6 weeks for people with hypertension leads to an increased risk of cardiovascular events
- Lowering blood pressure and reducing cardiovascular risk (1<sup>o</sup> and 2<sup>o</sup> prevention) is very effective at preventing heart attacks and strokes and premature death.
- UCLPartners Size of the Prize indicates the heart attacks and strokes that could be prevented by increasing the detection and management of high blood pressure.

[Search UCLP Size of the Prize for data on every ICB and PCN \(BP & Cholesterol\)](#)


# Adapting the Proactive Care Frameworks for Local Use

1. The UCLP Hypertension Framework supports practices and PCNs with search and stratification tools, pathways, resources and training:
  - To prioritise patients – who do we need to see now and who can we safely phase for later review?
  - To determine who has home BP monitors and support patients to buy valid monitors and submit accurate readings
  - To use the wider workforce to support patient education, self-management and lifestyle change
2. The UCLP Hypertension Framework will align with and support other local interventions for hypertension, e.g.:
  - Virtual group consultations to teach and check BP technique
  - Provision of free or loaned BP monitors to improve access
  - Targeting implementation to reduce health inequalities
  - Local quality improvement schemes for treatment optimisation



# Stratification and Management of High Blood Pressure

# Hypertension- Stratification and Management



\*QRISK 3 score is recommended to assess CV risk for patients with Severe Mental Illness, Rheumatoid Arthritis, Systemic Lupus Erythematosus, those taking antipsychotics or oral steroids; <sup>§</sup>ARRS = Additional Role Reimbursement Scheme

\*\* see slide 11

# Clinic vs home / ambulatory BP readings

Clinic BP reading **	Equivalent Home BP
BP = 180/120mmHg	BP = 170/115mmHg
BP = 160/100mmHg	BP = 150/95mmHg
BP = 150/90mmHg	BP = 145/85mmHg
BP = 140/90mmHg	BP = 135/85mmHg

# Lifestyle Modifications

Modification	Recommendation	Approximate Systolic Blood Pressure Reduction (mm Hg)**
Weight loss	Maintain normal body weight	5–20 per 10-kg weight loss
DASH-type diet*	Consume a diet rich in fruits, vegetables, and low-fat dairy products with reduced saturated and total fat	8–14
Reduced salt intake	Reduce daily dietary sodium intake	2–8
Physical activity	Regular aerobic physical activity (at least 30 min/day, most days of the week)	4–9
Moderation of alcohol intake	Limit consumption to 2 drinks/day in men and 1 drink/day in women and lighter-weight persons	2–4


**In monotherapy, most drugs achieve systolic BP reductions of ~ 10 to 15 mmHg**

[https://journals.lww.com/md-journal/Fulltext/2016/07260/Treatment\\_efficacy\\_of\\_anti\\_hypertensive\\_drugs\\_in.16.aspx](https://journals.lww.com/md-journal/Fulltext/2016/07260/Treatment_efficacy_of_anti_hypertensive_drugs_in.16.aspx)

\*DASH, Dietary Approaches to Stop Hypertension. Effects of implementing these modifications are time and dose dependent and could be greater for some patients.

\*\*Vooradi S, Mateti UV. A systemic review on lifestyle interventions to reduce blood pressure. J Health Res Rev [serial online] 2016 [cited 2021 Apr 27];3:1-5. Available from: [https://www.researchgate.net/figure/Reducing-blood-pressure-through-nonpharmacologic-measures-and-their-effects-and-target\\_tbl1\\_289571702](https://www.researchgate.net/figure/Reducing-blood-pressure-through-nonpharmacologic-measures-and-their-effects-and-target_tbl1_289571702)

# NICE Hypertension Treatment Pathway (NG136)



Use clinical judgement for people with frailty or multimorbidity

Offer lifestyle advice and continue to offer it periodically

**Monitoring treatment**

Use clinic BP to monitor treatment

Measure standing and sitting BP in people with:

- Type 2 diabetes or
- Symptoms of postural hypotension or
- Aged 80 and over

Advice people who want to self monitor to use HBPM. Provide training and advice

Consider AMPM or HBPM, in addition to clinic BP, for people with white-coat effect or masked hypertension

**BP targets**

Reduce and maintain BP to the following targets:

Age <80 years:

- Clinic BP <140/90 mmHg
- ABPM/HBPM <135/85mmHg

**Postural hypotension:**


- Base target on standing BP

**Frailty or multimorbidity:**

- Use clinical judgement


Pathway adapted from NICE Guidelines (NG136) Visual Summary  
<https://www.nice.org.uk/guidance/ng136/resources/visual-summary-pdf-6899919517>  
 Abbreviations: ACEi: ACE inhibitor, ARB: Angiotensin II Receptor Blocker, CCB: Calcium Channel Blocker, ABPM: Ambulatory Blood Pressure Monitoring, HBPM: Home Blood Pressure Monitoring

# Home Blood Pressure Monitoring Pathway



# Management of Broader Cardiovascular Risk in Hypertension:

## Detecting Atrial Fibrillation



# Detection and Management of AF in Patients with Hypertension

- Palpate pulse and if irregular or patient uncertain:
  - Assess for AF using ECG or [remote devices](#).
- If AF is confirmed, undertake stroke and bleeding risk assessment and anticoagulate as appropriate.
- Ensure following information is followed for an accurate blood pressure (BP) measurement:
  - Patients **without** AF:
    - Take 2 BP readings. If the values are more than 5mmHg apart, do a third BP reading and take an average of the lowest 2 BP readings.
  - Patients **with** AF:
    - Take blood pressure twice in the morning and twice in the evening for 4 consecutive days and then calculate an average of the values. (An average of 14 readings is required)

**Please refer to UCLP AF pathway for detailed guidance:**

<https://uclpartners.com/our-priorities/cardiovascular/proactive-care/cvd-resources/>



# **Management of Broader Cardiovascular Risk in Hypertension:**

## **Cholesterol Management**


# Managing High Cholesterol and Cardiovascular Risk in People with Hypertension

The following slides will help clinicians manage the broader cardiovascular risk in people with hypertension:

- **Pre-existing cardiovascular disease**
  - Optimise lifestyle
  - Use of high intensity statins at maximal appropriate dose
- **No pre-existing cardiovascular disease**
  - Optimise lifestyle and lipid lowering therapy as primary prevention in people with:
    - QRisk >10% in ten years
    - CKD 3-5
    - Type 1 Diabetes for >10 years or over age 40
- **All patients:**
  - Responding to possible statin intolerance
  - Managing muscle symptoms and abnormal LFTs in people taking statins
- **Please refer to UCLP lipid pathway for detailed guidance:**

<https://uclpartners.com/our-priorities/cardiovascular/proactive-care/cvd-resources/>


# Lipid Optimisation Pathway for Secondary Prevention<sup>1</sup>



**Lipid lowering therapy should be offered to all patients with established CVD<sup>1</sup>**

1. [NICE NG238: Cardiovascular disease: risk assessment and reduction, including lipid modification](#)
2. Dose may be limited, for example if:
  - CKD: eGFR<60ml/min – recommended starting dose - atorvastatin 20mg
  - Drug interactions
  - Drug intolerance
  - Older age / frailty
3. See [statin intensity table](#).
4. Use shared-decision making and incorporate patient preference in treatment and care decisions.
5. NICE Guidance PCSK9i(mAB): [Evolocumab](#), [Alirocumab](#)


# Optimisation Pathway for Patients with High Cardiovascular Risk – Primary Prevention<sup>1,2</sup>



Lipid lowering therapy should be offered to all patients at high CV risk<sup>2</sup>. (It may also be considered in individuals with QRisk < 10% )<sup>1</sup>  
[People living with HIV should be offered statin therapy regardless of QRisk score<sup>5</sup>](#)

1. [NICE NG238: Cardiovascular disease: risk assessment and reduction, including lipid modification](#)
2. High cardiovascular risk:
  - QRisk ≥ 10% in ten years
  - CKD: eGFR<60ml/min or albuminuria
  - Type 1 Diabetes for >10 years or over age 40, nephropathy or other CV risk factors
3. See [statin intensity table](#).
4. Use shared-decision making and incorporate patient preference in treatment and care decisions.
5. [BHIVA rapid guidance on the use of statins for primary prevention of cardiovascular disease in people living with HIV v2](#)

# Statin Initiation for Primary Prevention for People Living with HIV (PLWH) 40 years and above\*



# Common Statin/ARV Interactions and Recommended Doses for PLWH



ARV Regimen	Effect on Statins	Recommended atorvastatin starting dose	Maximum atorvastatin dose	Recommended rosuvastatin starting dose	Maximum rosuvastatin dose
<b>Ritonavir- or cobicistat-boosted darunavir</b>	Increased atorvastatin and rosuvastatin concentrations.	10mg	40mg	5mg	20mg
<b>Ritonavir- or cobicistat-boosted elvitegravir</b>	Increased atorvastatin and rosuvastatin concentrations.	10mg	40mg	5mg	20mg
<b>Ritonavir- or cobicistat-boosted atazanavir</b>	Significantly higher atorvastatin and rosuvastatin levels.	10mg	10mg	5mg	10mg
<b>Lopinavir/ritonavir</b>	Significantly higher atorvastatin and rosuvastatin levels.	10mg	20mg	5mg	10mg
<b>Efavirenz</b>	Variable reductions in atorvastatin. Rosuvastatin preferred first line.	20mg	80mg	10mg	40mg
<b>Other ARV regimens</b>	See resources below or seek advice from a lipid / HIV specialist				

Standard dosing of Ezetimibe is advised for all ARV regimens

Please note some antiretrovirals, such as boosted protease inhibitors (e.g. darunavir/ritonavir/cobicistat/atazanavir) and efavirenz can increase lipids, while others are more lipid-friendly. Consider referring patients with persistently elevated lipids to their HIV clinic for optimisation of their antiretroviral regimen.

For further information and advice about interactions: <https://www.hiv-druginteractions.org/checker>



# Statin Intolerance Pathway

## Important considerations

- Most adverse events attributed to statins are no more common than placebo<sup>1</sup>
- Consider food and drug interactions which may be contributing to adverse effects – see Summary of Product Characteristics (SmPC)<sup>2,3</sup>
- Stopping statin therapy is associated with an increased risk of major CV events. It is important not to label patients as ‘statin intolerant’ without structured assessment
- If a person is not able to tolerate a high-intensity statin, aim to treat with the maximum tolerated dose
- A statin at any dose reduces CVD risk – consider annual review for patients not taking statins to review cardiovascular risk and interventions

## A structured approach to reported adverse effects of statins

- Stop for 4-6 weeks.
- If symptoms persist, they are unlikely to be due to statin
- Restart and consider lower initial dose
- If symptoms recur, consider trial with alternative statin
- If symptoms persist, consider [ezetimibe](#) +/- [bempedoic acid](#)

1. (Collins et al systematic review, Lancet 2016)
2. SmPC: Atorvastatin  
<https://www.medicines.org.uk/emc/product/5274/smpc#gref>
3. SmPC: Rosuvastatin  
<https://www.medicines.org.uk/emc/product/4366/smpc#gref>



# Resources

# For Local Decision: Options for Purchasing Home BP Machines

## Validated devices

- A list of validated devices for home use can be found at:  
<https://giftshop.bhf.org.uk/health/blood-pressure-monitors>. Validated devices for home use are accurate for up to 5 years after purchase

(Hodgkinson JA et al. 2020 Accuracy of blood-pressure monitors owned by patients with hypertension (ACCU-RATE study): a cross-sectional, observational study in central England. BJGP 1 June 2020; bjgp20X710381. DOI: <https://doi.org/10.3399/bjgp20X710381>)

## Considerations

- Upper arm blood pressure devices preferred
- Basic model (~£20) is suitable for most patients
- Ensure patient has the correct cuff size based on arm circumference
- Bluetooth connectivity allows automatic transfer of data into a patient held device. However few NHS services are able to interface with these data portals at this time and Bluetooth enabled devices are more expensive to purchase

# Resources for Remote Diagnostics and Monitoring

## **Newly identified irregular heart rhythm in people with high blood pressure**

- Fibrichck (needs smartphone) [www.fibrichck.com/](http://www.fibrichck.com/) and ask patient to monitor morning and evening for 7 days
- Utilise mobile ECG technology, if available e.g.:
  - Kardia by AliveCor (needs smartphone): [www.alivecor.co.uk/kardiamobile](http://www.alivecor.co.uk/kardiamobile)
  - Zenicor: <https://zenicor.com/>

## **ACR - home urine testing**

- Healthy.io <https://healthy.io/urinalysis-products/>

# Resources for Patients

## Resources on high blood pressure and how to manage it:

- **British Heart Foundation** hub for managing blood pressure at home aimed at patients  
[www.bhf.org.uk/bloodpressureathome](http://www.bhf.org.uk/bloodpressureathome)
- **Stroke Association:** [www.stroke.org.uk/what-is-stroke/are-you-at-risk-of-stroke/high-blood-pressure](http://www.stroke.org.uk/what-is-stroke/are-you-at-risk-of-stroke/high-blood-pressure)

## Monitoring your blood pressure at home:

- [How to check your blood pressure using a blood pressure machine \(video\)](#)
- [Tips for measuring your BP at home](#)
- [Home monitoring diary for patients](#)
- [Validated BP monitors for home use](#)
- [How to choose a BP monitor](#)
- [Remote Blood Pressure Monitoring Video for Patients \(English\)](#)
- [Remote Blood Pressure Monitoring Video for Patients \(Other languages\)](#)

# Resources for Patients

## How to assess pulse rhythm at home

British Heart Foundation: [How to take your pulse video](#)

Heart Rhythm Alliance: [Know Your Pulse Factsheet](#), [What is an Arrhythmia?](#)

## Diet

Heart UK: [Blood fats explained](#)

Providing information on healthy eating from the [NHS website](#)

Advice and guidance on losing weight including useful apps and healthy recipes on the [Better Health website](#)

[NHS advice on lowering cholesterol levels](#)

## Smoking cessation

[NHS support](#), stop smoking aids, tools and practical tips

## Exercise

[Getting active around the home](#): tips, advice and guidance on how to keep or get active in and around the home from Sport England

[Dance to health](#): Online dance programme especially tailored to people over 55 years old

## Alcohol

[Heart UK alcohol guidance](#) & [NHS Drink Less guidance](#)



# Implementation Support

# Implementation & Support Package

Implementation Support is critical to enable sustainable and consistent spread. UCLPartners has developed a support package for the Integrated Care Systems within our geography covering the following components. The resources below can be accessed via the UCLP website: [Proactive care frameworks – UCLPartners](#).

UCLPartners is one of 15 [Health Innovation Networks](#) (HINs) across England and all 15 have a priority around CVD. Please reach out to your local HIN to understand what support they might be able to provide. Please note each varies in its approach and offer.

## Search and stratify

**Comprehensive search tools** for EMIS and SystmOne to stratify patients

- Pre-recorded webinar as to how to use the searches.
- Online FAQs to troubleshoot challenges with delivery of the search tools.

## Workforce training and support

**Training tailored to each staff grouping (e.g. some ARRS\* roles) and level of experience**

- **Delivery:** Scripts provided as well as training on how to use these underpinned with motivational interviewing/ health coaching training to enable adult-to-adult conversations.
- **Practical support:** [Recommended training](#) e.g. correct inhaler technique; correct BP technique, Very Brief Advice for smoking cessation, physical activity etc.
- **Digital implementation** support: how to get patients set up with appropriate digital.
- **Education** sessions on conditions.
- **Communities of Practice.**

## Digital support tools

**Digital resources** to support remote management and self-management in each condition.

**Implementation** toolkits available where required, e.g. MyCOPD.

Support available from UCLP's commercial and innovation team for implementation.

# Thank you

Sign up to our monthly newsletter to receive the latest news, opportunities and events from UCLPartners



[UCLPartners.com/newsletter](https://UCLPartners.com/newsletter)

---

For any enquiries, please contact us via email:

[\*\*primarycare@uclpartners.com\*\*](mailto:primarycare@uclpartners.com)



[www.uclpartners.com](https://www.uclpartners.com)



[@uclpartners](https://twitter.com/uclpartners)



[linkedin/company/uclpartners](https://linkedin/company/uclpartners)



[comms@uclpartners.com](mailto:comms@uclpartners.com)



# Version Tracker

Version	Edition	Changes Made	Date amended	Review due
2	2.0	<ul style="list-style-type: none"> <li>Incorporated lipid management content for patients with multi-morbidity</li> <li>Added lifestyle interventions for patient self-management</li> <li>Added statin intensity table for reference</li> </ul>		
3	3.0	<ul style="list-style-type: none"> <li>Added option of bempedoic acid</li> <li>Amended slide on managing high cholesterol</li> <li>Amended slide on managing/detecting AF</li> </ul>	August 2021	February 2022
4	4.0	<ul style="list-style-type: none"> <li>Included Inclisiran into lipid management pathway</li> <li>Updated priority groups</li> <li>Updated web links for resources</li> </ul>	July 2022	July 2023
5	5.0	<ul style="list-style-type: none"> <li>Introduction slides updated</li> <li>HCA roles amended to ARRS roles</li> <li>Lipid pathway treatment targets updated to align with NICE and AAC guidance</li> </ul>	December 2022	December 2023
5	5.1	<ul style="list-style-type: none"> <li>Updated cholesterol pathway slides</li> <li>Amended introduction slides</li> <li>Added link for home BP monitoring video in other languages</li> </ul>	September 2023	September 2024
6	6.0	<ul style="list-style-type: none"> <li>Updated cholesterol pathway slides to align with NICE</li> <li>Updated implementation slide</li> </ul>	January 2024	January 2025
6	6.1	<ul style="list-style-type: none"> <li>Slide set moved onto new UCLP branding</li> <li>Size of the prize slide added</li> </ul>	April 2024	April 2025
7	7.0	<ul style="list-style-type: none"> <li>Secondary care pathway slide amended to include statin monitoring info and high intensity statin examples incorporated within pathway.</li> </ul>	June 2024	January 2025
8	8.0	<ul style="list-style-type: none"> <li>Updated in line with NICE guidance</li> <li>Revised searches to align with NICE BP thresholds</li> <li>Addition of lipid management in people living with HIV</li> </ul>	April 2025	April 2026
9	9.0	<ul style="list-style-type: none"> <li>Updated lipid optimisation pathway for secondary prevention</li> </ul>	Feb 2026	Feb 2027